

A NOTE ON NON-AMENABILITY OF $\mathcal{B}(\ell_p)$ FOR $p = 1, 2$

NARUTAKA OZAWA

ABSTRACT. This is an expository note on non-amenability of the Banach algebra $\mathcal{B}(\ell_p)$ for $p = 1, 2$. These were proved respectively by Connes and Read via very different methods. We give a single proof which reproves both.

1. INTRODUCTION

Johnson [Jo1] introduced the notion of amenability for Banach algebra and asked whether $\mathcal{B}(X)$, the Banach algebra of bounded linear map on a Banach space X , can be amenable. This problem is still widely open. For the special case where $X = \ell_p$, it was proved by Connes [Co] that $\mathcal{B}(\ell_2)$ and $\prod_{n=1}^{\infty} \mathcal{B}(\ell_2^n)$ are not amenable, and more recently, by Read [Re] that $\mathcal{B}(\ell_1)$ and $\prod_{n=1}^{\infty} \mathcal{B}(\ell_p^n)$ for $p \in [1, \infty] \setminus \{2\}$ are not amenable. It is left open whether $\mathcal{B}(\ell_p)$ is amenable or not for $p \in (1, \infty) \setminus \{2\}$. A somewhat simplified proof of Read's results was given by Pisier [Pi2] where it was observed that the proof also gives the non-amenability of 'regular' $\mathcal{B}(\ell_p)$. In this note, we give a single proof which reproves all above results. The proof is inspired from Read's approach and was suggested by Pisier.

Theorem 1.1. *The following Banach algebras are not amenable; $\mathcal{B}(\ell_p)$ for $p \in \{1, 2, \infty\}$, $\prod_{n=1}^{\infty} \mathcal{B}(\ell_p^n)$ for $p \in [1, \infty]$ and $\mathcal{B}_r(\ell_p)$ for $p \in [1, \infty]$.*

Here, $\mathcal{B}_r(\ell_p)$ is the Banach algebra consisting of those operators $x \in \mathcal{B}(\ell_p)$ with

$$\|x\|_{\mathcal{B}_r(\ell_p)} := \| |x| \|_{\mathcal{B}(\ell_p)} < \infty,$$

where $|x| = (|x_{ij}|)_{i,j}$ for $x = (x_{ij}) \in \mathcal{B}(\ell_p)$. We use the following definition of amenability of Banach algebras, which is equivalent to the standard one by Johnson's theorem [Jo2]. A general reference for amenable Banach algebra is [Ru].

Definition 1.2. A unital Banach algebra A is *amenable* if there exists a constant $C > 0$ with the following property; for any finite set $F \subset A$ and $\varepsilon > 0$, there exists $T = \sum_{i=1}^r a_i \otimes b_i \in A \otimes A$ such that

- (i) $\text{prod}(T) = \sum_{i=1}^r a_i b_i = 1$,
- (ii) $\|x \cdot T - T \cdot x\|_{\wedge} = \|\sum_{i=1}^r x a_i \otimes b_i - \sum_{i=1}^r a_i \otimes b_i x\|_{\wedge} \leq \varepsilon$ for $x \in F$,
- (iii) $\|T\|_{\wedge} \leq C$,

where $\|\cdot\|_{\wedge}$ denotes the projective tensor norm on $A \otimes A$.

It turns out that the condition (iii) is irrelevant to Theorem 1.1.

2000 *Mathematics Subject Classification.* Primary 46H20; Secondary 46B07.

2. SOME FACTS ON ℓ_p

In this section, we prove some elementary facts on ℓ_p . Fix $p \in [1, \infty]$ and let q be so that $1/p + 1/q = 1$. We denote by $\{\delta_m\}$ (resp. $\{\delta_m^*\}$) the standard basis of ℓ_p (resp. ℓ_q). We temporarily denote by $\|x\|_{\text{reg}} = \| |x| \|$ for x in $\mathcal{B}(\ell_p)$ or $\mathcal{B}(\ell_q)$.

Lemma 2.1. *For $x \in \mathcal{B}(\ell_p, \ell_p^N)$ and $y \in \mathcal{B}(\ell_q, \ell_q^N)$, we have*

- (i) $\sum_{m=1}^{\infty} \|x\delta_m\|_q \|y\delta_m^*\|_p \leq N \|x\| \|y\|,$
- (ii) $\sum_{m=1}^{\infty} \|x\delta_m\|_p \|y\delta_m^*\|_q \leq N \|x\|_{\text{reg}} \|y\|_{\text{reg}},$
- (iii) $\sum_{m=1}^{\infty} \|x\delta_m\|_2 \|y\delta_m^*\|_2 \leq N (\|x\| \|x\|_{\text{reg}} \|y\| \|y\|_{\text{reg}})^{1/2}.$

Remark 2.2. The estimate $\sum_{m=1}^{\infty} \|x\delta_m\|_2 \|y\delta_m^*\|_2 \leq N \|x\| \|y\|$ needs not be true (even up to constant multiple) for $x \in \mathcal{B}(\ell_p, \ell_p^N)$ and $y \in \mathcal{B}(\ell_q, \ell_q^N)$. However, it is not hard to see (from the following proof) this is true if $p \in \{1, 2\}$ or if $x \in \mathcal{B}(\ell_p^N)$ and $y \in \mathcal{B}(\ell_q^N)$.

Proof. The part (i) follows from Hölder inequality;

$$\begin{aligned} \sum_{m=1}^{\infty} \|x\delta_m\|_q \|y\delta_m^*\|_p &\leq \left(\sum_{m=1}^{\infty} \|x\delta_m\|_q^q \right)^{1/q} \left(\sum_{m=1}^{\infty} \|y\delta_m^*\|_p^p \right)^{1/p} \\ &= \left(\sum_{k=1}^N \|x^* \delta_k^*\|_q^q \right)^{1/q} \left(\sum_{k=1}^N \|y^* \delta_k\|_p^p \right)^{1/p} \\ &\leq N \|x\| \|y\|. \end{aligned}$$

For the part (ii), we may assume $x = |x|$. Then, by the duality $\ell_q = (\ell_p)^*$, we have

$$\begin{aligned} \left(\sum_{m=1}^{\infty} \|x\delta_m\|_p^q \right)^{1/q} &= \sup_{\lambda} \sum_{m=1}^{\infty} \lambda_m \|x\delta_m\|_p \leq \sup_{\lambda} \sum_{m=1}^{\infty} \lambda_m \|x\delta_m\|_1 \\ &= \sup_{\lambda} \|x\lambda\|_1 \leq N^{1/q} \sup_{\lambda} \|x\lambda\|_p \\ &\leq N^{1/q} \|x\|_{\text{reg}}, \end{aligned}$$

where the supremum is taken over all $\lambda = (\lambda_m)_m$ with $\|\lambda\|_p \leq 1$. The rest of the proof of part (ii) is same as (i). Finally, the part (iii) follows from (i) and (ii) combined with the Cauchy-Schwarz inequality and the fact $\|\lambda\|_2 \leq (\|\lambda\|_p \|\lambda\|_q)^{1/2}$. $\square$

3. SOME FACTS ON $\text{SL}(3, \mathbb{Z})$

In this section, we collect some known facts on the group $\text{SL}(3, \mathbb{Z})$. For notational simplicity, we denote $\text{SL}(3, \mathbb{Z})$ by Γ . The group Γ has Kazhdan's property (T). In particular, it is finitely generated. (In fact, it is generated by two elements.) A general reference for Kazhdan's property (T) is [HV].

Theorem 3.1. *The group $\Gamma = \text{SL}(3, \mathbb{Z})$ has Kazhdan's property (T). Namely, for any finite set of generators $\mathcal{S}$ of Γ , there exists a positive number $R = R(\Gamma, \mathcal{S})$ with*

the following property; for any unitary representation Γ on a Hilbert space $\mathcal{H}$ and a vector $\xi \in \mathcal{H}$, there exists a π -invariant vector $\eta \in \mathcal{H}$ such that

$$\|\xi - \eta\| \leq R \max_{g \in \mathcal{S}} \|\pi(g)\xi - \xi\|.$$

This theorem is non-trivial only when $\max_{g \in \mathcal{S}} \|\pi(g)\xi - \xi\| < \|\xi\|/R$ since the zero vector is π -invariant. See Appendix for a related fact on actions on Banach spaces.

We next present explicit examples of actions of Γ . For each prime number l , let $\mathbb{F}_l = \mathbb{Z}/l\mathbb{Z}$ be the finite field with l elements and define the “finite projective plane” Λ_l by $\Lambda_l = (\mathbb{F}_l^3 - \{0\})/\mathbb{F}_l^\times$. The action Γ on $\mathbb{F}_l^3$ (through linear transformation by $\text{SL}(3, \mathbb{F}_l)$) induces an action σ_l on the set Λ_l . It is not hard to see that σ_l is 2-transitive, i.e., for any $s_{ij} \in \Lambda_l$ ($i, j \in \{1, 2\}$) with $s_{1j} \neq s_{2j}$ for $j \in \{1, 2\}$, there exists $g \in \Gamma$ such that $\sigma_l(g)(s_{i1}) = s_{i2}$ for $i = 1, 2$. In other words, the product action $\sigma_l \times \sigma_l$ of Γ on $\Lambda_l \times \Lambda_l$ has exactly two orbits (diagonals and off-diagonals). For $p \in [1, \infty]$, we denote by π_l^p the isometric action of Γ on $\ell_p(\Lambda_l)$ induced from σ_l . Let $\Lambda = \bigsqcup_l \text{prime } \Lambda_l$ be the disjoint union of Λ_l ’s and let $\pi^p = \bigoplus \pi_l^p$ be the isometric action of Γ on $\ell_p(\Lambda)$. For a later use, we choose a subset $C_l \subset \Lambda_l$ with $|C_l| = (l^2 + l)/2 = (|\Lambda_l| - 1)/2$ and let $v_l^p \in \mathcal{B}(\ell_p(\Lambda_l))$ be the isometry given by $v_l^p \delta_s = \delta_s$ if $s \in C_l$ and $v_l^p \delta_s = -\delta_s$ if $s \notin C_l$. Finally, set $v^p = \bigoplus v_l^p \in \mathcal{B}(\ell_p(\Lambda))$.

4. SOME FACTS ON $\ell_2 \hat{\otimes} \ell_2$

In this section, we review a well-known fact about operator inequality. A general reference is [Bh]. Let S_1 (resp. S_2) be the Banach space of trace (resp. Hilbert-Schmidt) class operators on ℓ_2 . The ‘noncommutative’ Mazur map $\varphi: S_1 \rightarrow S_2$ is defined by $\varphi(T) = U|T|^{1/2}$, where $T = U|T|$ is the polar decomposition and modulus and square root are taken as an operator on a Hilbert space. By the uniqueness of polar decomposition, we have $\varphi(USU^*) = U\varphi(S)U^*$ for any unitary operator U . As in the case of ordinary Mazur map, we have the following.

Theorem 4.1. *The noncommutative Mazur map $\varphi: S_1 \rightarrow S_2$ is a uniform homeomorphism between the unit ball of S_1 and that of S_2 .*

Proof. We give a proof for the reader’s convenience. Temporally, we denote by $\|\cdot\|_p$ the Schatten p -norm. It suffices to show that φ is a uniform homeomorphism between the spheres. Let $S = U|S|$ and $T = V|T|$ be given so that $\|S\|_1 = 1 = \|T\|_1$. Let $\varepsilon = \|S - T\|_1$. Then, $\||S| - |T|\|_1 \leq 2\varepsilon^{1/2}$ by [Ko] and $\||S|^{1/2} - |T|^{1/2}\|_2 \leq 2\varepsilon^{1/4}$ by [PS]. It follows that

$$\begin{aligned} \||S| - V^*U|S|^{1/2}|T|^{1/2}\|_1 &\leq \||S| - V^*S\|_1 + \|V^*U|S|^{1/2}\|_2 \||S|^{1/2} - |T|^{1/2}\|_2 \\ &\leq \|S - T\|_1 + \||S| - |T|\|_1 + 2\varepsilon^{1/4} \\ &\leq \varepsilon + 2\varepsilon^{1/2} + 2\varepsilon^{1/4}. \end{aligned}$$

Therefore, we have

$$\|\varphi(S) - \varphi(T)\|_2^2 = 2 - 2\Re \operatorname{Tr}(\varphi(T)^* \varphi(S)) \leq 2\varepsilon + 4\varepsilon^{1/2} + 4\varepsilon^{1/4}.$$

This proves the uniform continuity of φ . The uniform continuity of φ^{-1} follows from the Cauchy-Schwarz inequality. $\square$

Let us identify S_1 (resp. S_2) with the projective (resp. ℓ_2 -) tensor product $\ell_2 \hat{\otimes} \ell_2$ (resp. $\ell_2 \otimes_2 \ell_2$). The identification is given by $\xi \otimes \eta \mapsto T_{\xi, \eta}$, where $T_{\xi, \eta}(\zeta) = (\zeta, \bar{\xi})\eta$. (The symbol $\bar{\xi}$ means the complex conjugate of $\xi \in \ell_2$.) Then, the noncommutative Mazur map $\varphi: \ell_2 \hat{\otimes} \ell_2 \rightarrow \ell_2 \otimes_2 \ell_2$ satisfies $\varphi \circ (U \otimes \bar{U}) = (U \otimes \bar{U}) \circ \varphi$ for any unitary operator U and is uniformly homeomorphic on the spheres.

5. PROOF OF NON-AMENABILITY

Proof. Let A be one of the Banach algebras listed in Theorem 1.1. Set q be so that $1/p + 1/q = 1$. To give a finite set $F \subset A$ which violates the definition of amenability, we fix a finite set $\mathcal{S}$ of generators of Γ . For notational simplicity, we set $\mathcal{S}^+ = \mathcal{S} \cup \{g_v\}$ and $\pi^p(g_v) = v^p$. Then,

$$F = \{\pi^p(g) : g \in \mathcal{S}^+\} \subset \prod \mathcal{B}(\ell_p(\Lambda_l)) \subset \mathcal{B}(\ell_p(\Lambda))$$

(cf. Section 3) is a finite subset in A . We choose $\varepsilon > 0$ later. Suppose there exists $T = \sum_{i=1}^r a_i \otimes b_i \in A \otimes A$ which satisfy the conditions (i) and (ii) in Definition 1.2.

We fix a prime number l until the very last step. For each $m \in \Lambda$, we define $T_l(m) \in \ell_2(\Lambda_l) \hat{\otimes} \ell_2(\Lambda_l)$ by

$$T_l(m) = \sum_{i=1}^r P_l a_i \delta_m \otimes P_l^* b_i^* \delta_m^* \in \ell_2(\Lambda_l) \hat{\otimes} \ell_2(\Lambda_l),$$

where P_l is the (orthogonal) projection from $\ell_p(\Lambda)$ onto $\ell_p(\Lambda_l)$. By Lemma 2.1 (and the following remark), we have $\sum_{m \in \Lambda} \|T_l(m)\|_{\ell_2(\Lambda_l) \hat{\otimes} \ell_2(\Lambda_l)} \leq |\Lambda_l| \|T\|_{A \hat{\otimes} A}$. Similarly

$$\begin{aligned} \sum_{m \in \Lambda} \|T_l(m) - (\pi_l^2(g) \otimes \pi_l^2(g))T_l(m)\|_{\ell_2(\Lambda_l) \hat{\otimes} \ell_2(\Lambda_l)} \\ \leq |\Lambda_l| \|T - \pi^p(g) \cdot T \cdot \pi^p(g)^{-1}\|_{A \hat{\otimes} A} \leq |\Lambda_l| \varepsilon \end{aligned}$$

for every $g \in \mathcal{S}^+$. On the other hand, since $\sum_{i=1}^r a_i b_i = 1$, we have

$$\sum_{m \in \Lambda} \|T_l(m)\|_{\ell_2(\Lambda_l) \hat{\otimes} \ell_2(\Lambda_l)} \geq \sum_{m \in \Lambda} \langle P_l a_i \delta_m, P_l^* b_i^* \delta_m^* \rangle = \operatorname{Tr}(P_l) = |\Lambda_l|.$$

Therefore, there exists $m_l \in \Lambda$ such that

$$\|T_l(m_l) - (\pi_l^2(g) \otimes \pi_l^2(g))T_l(m_l)\|_{\ell_2(\Lambda_l) \hat{\otimes} \ell_2(\Lambda_l)} \leq \varepsilon |\mathcal{S}^+| \|T_l(m_l)\|_{\ell_2(\Lambda_l) \hat{\otimes} \ell_2(\Lambda_l)}$$

for all $g \in \mathcal{S}^+$. Set $S_l = T_l(m_l) / \|T_l(m_l)\|_{\wedge}$. The application of the noncommutative Mazur map φ to this inequality yields (cf. the last remark in Section 4) that

$$\|\varphi(S_l) - (\pi_l^2(g) \otimes \pi_l^2(g))\varphi(S_l)\|_{\ell_2(\Lambda_l) \otimes_2 \ell_2(\Lambda_l)} \leq \omega_\varphi(\varepsilon |\mathcal{S}^+|) =: \delta_0$$

for all $g \in \mathcal{S}^+$, where ω_φ is the modulus of continuity of φ (cf. Appendix). Now, since $\pi_l^2 \otimes \pi_l^2$ on $\ell_2(\Lambda_l) \otimes \ell_2(\Lambda_l)$ is a unitary representation of Γ , Theorem 3.1 is applicable. It follows that there exists a $(\pi_l^2 \otimes \pi_l^2)$ -invariant vector X such that

$$\|\varphi(S_l) - X\|_{\ell_2(\Lambda_l) \otimes \ell_2(\Lambda_l)} \leq R\delta_0,$$

where $R = R(\Gamma, \mathcal{S})$ is the constant appearing in Theorem 3.1. But since Γ action on Λ_l is 2-transitive, $X = \lambda I + \mu E$ for some $\lambda, \mu \in \mathbb{C}$, where $I = |\Lambda_l|^{-1/2} \sum_{s \in \Lambda_l} \delta_s \otimes \delta_s$ and $E = |\Lambda_l|^{-1} \sum_{s, t \in \Lambda_l} \delta_s \otimes \delta_t$. It follows that

$$\begin{aligned} (2 - 2|\Lambda_l|^{-2})^{1/2} |\mu| &= |\mu| \|E - (v^2 \otimes v^2)E\|_{\ell_2(\Lambda_l) \otimes \ell_2(\Lambda_l)} \\ &= \|X - (v^2 \otimes v^2)X\|_{\ell_2(\Lambda_l) \otimes \ell_2(\Lambda_l)} \\ &\leq 2R\delta_0 + \|\varphi(S_l) - (v^2 \otimes v^2)\varphi(S_l)\|_{\ell_2(\Lambda_l) \otimes \ell_2(\Lambda_l)} \\ &\leq (2R + 1)\delta_0. \end{aligned}$$

Therefore,

$$\|\varphi(S_l) - \lambda I\|_{\ell_2(\Lambda_l) \otimes \ell_2(\Lambda_l)} \leq (3R + 1)\delta_0 =: \delta_1.$$

and in particular $|\lambda| \geq \|\varphi(S_l)\| - \delta_1 = 1 - \delta_1$. On the other hand, since $\varphi(S_l)$ is of rank $\leq r$ (as a Hilbert-Schmidt operator on $\ell_2(\Lambda_l)$), we have

$$(1 - \frac{r}{|\Lambda_l|})|\lambda|^2 \leq \|\varphi(S_l) - \lambda I\|_{\ell_2(\Lambda_l) \otimes \ell_2(\Lambda_l)}^2 = \delta_1^2.$$

Consequently,

$$r \geq (1 - \frac{\delta_1^2}{(1 - \delta_1)^2})|\Lambda_l| =: (1 - \delta(\varepsilon))|\Lambda_l|,$$

where $\delta: \mathbb{R}_+ \rightarrow [0, 1]$ is a function which does not depend on l (but depends on $\Gamma, \mathcal{S}$ and ω_φ) such that $\lim_{\varepsilon \rightarrow 0} \delta(\varepsilon) = 0$. Since l can be arbitrarily large, this is absurd when $\varepsilon > 0$ is chosen sufficiently small so that $\delta(\varepsilon) < 1$. $\square$

Remark 5.1. In the proof, one may use Corollary A.3, instead of Theorem 3.1, to deduce that $S_l \in \ell_2(\Lambda_l) \hat{\otimes} \ell_2(\Lambda_l)$ is close to a $(\pi_l^2 \otimes \pi_l^2)$ -invariant vector. The first half of the proof goes *mutatis mutandis* for $\mathcal{B}(\ell_p)$ and yields $S_l \in \ell_q(\Lambda_l) \hat{\otimes} \ell_p(\Lambda_l)$ which is almost $(\pi_l^q \otimes \pi_l^p)$ -invariant. However, since we do not know whether the unit ball of $\ell_q \hat{\otimes} \ell_p$ is uniformly embeddable into a Hilbert space, it is not clear whether Corollary A.3 is applicable in this case or not. We note that it is not even known whether $\ell_q \hat{\otimes} \ell_p$ has non-trivial cotype or not (cf. [Pi1]).

APPENDIX A. NON-UNIFORM EMBEDDABILITY OF EXPANDERS

In this appendix, we prove a result about embedding of expanders into a Banach space. Throughout this appendix, we denote by G a graph which is simple, connected and k -regular. We abuse the notation and identify G with its vertices. Further, we regard G as the metric space equipped with the graph distance. For $A \subset G$, the boundary ∂A is the set of (unoriented) edges which connects A to $G \setminus A$. Let us recall

some basic facts about expanders. A general reference for expanders is [Lu]. The expanding constant $h(G)$ of G is given by

$$h(G) = \inf\left\{\frac{|\partial A|}{|A|} : A \subset V \text{ with } |A| \leq |V|/2\right\}.$$

For $h > 0$, we say G is an (k, h) -expander if it is k -regular and $h(G) \geq h$.

Let M_1 and M_2 be metric spaces and let $\varphi: M_1 \rightarrow M_2$ be a map. The Lipschitz norm of φ is given by $\text{Lip}(\varphi) = \sup\{\text{dist}(\varphi(a), \varphi(b))/\text{dist}(a, b) : a \neq b\}$. When $M_1 = G$ is a graph, then

$$\text{Lip}(\varphi) = \sup\{\text{dist}(\varphi(s), \varphi(t)) : s, t \in G \text{ are adjacent}\}.$$

The modulus of continuity of φ is the function $\omega_\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ given by

$$\omega_\varphi(t) = \sup\{\text{dist}(\varphi(a), \varphi(b)) : a, b \in M_1 \text{ and } \text{dist}(a, b) \leq t\}.$$

The map φ is said to be uniformly continuous if $\lim_{t \rightarrow 0} \omega_\varphi(t) = 0$. The metric space M_1 is said to be uniformly homeomorphic to M_2 if there is a bijection $\varphi: M_1 \rightarrow M_2$ such that both φ and φ^{-1} are uniformly continuous. Finally, M_1 is said to be uniformly embeddable into M_2 if M_1 is uniformly homeomorphic to a subset of M_2 .

We have the following concentration theorem for expanders embedded in a Banach space. A better estimate is known [LLR][Ma] for $X = \ell_p$ with $p \in [1, \infty)$.

Theorem A.1. *Let X be a Banach space whose unit ball is uniformly embeddable into a Hilbert space. Then, for any $k \in \mathbb{N}$ and $h > 0$, there exists a positive number $R = R(k, h, X)$ which satisfies the following; for any map f from a (k, h) -expander G into X , we have*

$$\frac{1}{|G|} \sum_{s \in G} \|f(s) - m\| \leq R \text{Lip}(f),$$

where $m = |G|^{-1} \sum_{s \in G} f(s)$ is the mean of f .

Any n -point metric space is isometric to a subset of ℓ_∞^n . A Banach space is said to have non-trivial cotype if it does not contain ℓ_∞^n 's uniformly. Thus a Banach space satisfying the conclusion of the theorem has non-trivial cotype (compare this with [En]). It would be interesting to know whether the converse is also true or not. We note that, by Odell and Schlumprecht's theorem [OS][Ch], a Banach space X with an unconditional basis and of non-trivial cotype has the unit ball which is uniformly homeomorphic to the unit ball of a Hilbert space. A general reference for related facts is [BL].

We apply the above theorem to the Caley graphs of a finite group. Recall that the Caley graph $G(\Delta, \mathcal{S})$ of a group Δ with a distinguished set of generators $\mathcal{S}$ is the graph whose vertices are elements of Δ and whose edges are pairs $\{g, h\}$ with $g^{-1}h \in \mathcal{S}$. The following is Margulis' celebrated construction of expanders (cf. [Lu]).

Theorem A.2. *Let Γ be a group with Kazhdan's property (T) and $\mathcal{S}$ be a finite set of generators of Γ . Then, there exists $h = h(\Gamma, \mathcal{S}) > 0$ such that for any finite quotient group Δ of Γ , the Cayley graph $G(\Delta, \mathcal{S})$ is an $(|\mathcal{S}|, h)$ -expander.*

Corollary A.3. *Let Γ be a group with Kazhdan's property (T), $\mathcal{S}$ be a finite set of generators of Γ and let X be a Banach space whose unit ball is uniformly embeddable into a Hilbert space. Then, there exists $R = R(\Gamma, \mathcal{S}, X) > 0$ which satisfies the following; for any representation $\pi: \Gamma \rightarrow \mathcal{B}(X)$ with a finite image and any $\xi \in X$, there exists a π -invariant vector η such that*

$$\|\xi - \eta\| \leq R \max_{g \in \Gamma} \|\pi(g)\|^2 \max_{h \in \mathcal{S}} \|\pi(h)\xi - \xi\|.$$

Proof. Apply Theorems A.1 and A.2 to the map $f: \pi(\Gamma) \ni \pi(g) \mapsto \pi(g)\xi \in X$. $\square$

The following lemma is well-known, but we include the proof for completeness.

Lemma A.4. *Let G be a (k, h) -expander. Then, we have the following.*

(i) *For any map $f: G \rightarrow \ell_1$, we have*

$$\frac{1}{|G|} \sum_{s \in G} \|f(s) - m\| \leq 2 \frac{k}{h} \text{Lip}(f),$$

where $m = |G|^{-1} \sum_{s \in G} f(s)$ is the mean of f .

(ii) *For any function $f: G \rightarrow \mathbb{R}_+$ with $|\{s \in G : f(s) \leq R_0\}| \geq |G|/2$, we have*

$$\frac{1}{|G|} \sum_{s \in G} f(s) \leq R_0 + \frac{k}{2h} \text{Lip}(f).$$

Proof. First, consider a function $g: G \rightarrow \mathbb{R}_+$ with $|\text{supp}(g)| \leq |G|/2$. Let E be the set of unoriented edges of G and let $A_r = \{s \in G : g(s) \geq r\}$. We note that $|A_r| \leq |\text{supp}(g)| \leq |G|/2$ and hence $|\partial A_r| \geq h|A_r|$ for $r > 0$. Since $|g(s) - g(t)|$ is the Lebesgue measure of the interval between $g(s)$ and $g(t)$, we have

$$\sum_{\{s, t\} \in E} |g(s) - g(t)| = \int_0^\infty |\partial A_r| \, dr \geq \int_0^\infty h|A_r| \, dr = h \sum_{s \in G} g(s).$$

Now the assertion (i) follows by considering each coordinate of ℓ_1 and applying the above inequality to the positive (or perhaps negative) part of $\Re f$ (resp. $\Im f$). The part (ii) follows from the above inequality for $g = \max\{0, f - R_0\}$. $\square$

Now, we give the proof of Theorem A.1.

Proof of Theorem A.1. We note that the Hilbert space ℓ_2 is uniformly embeddable into ℓ_1 (cf. [BL]). Fix a uniform embedding φ of the unit ball of X into ℓ_1 . Take $R > 0$ large enough so that it satisfies

$$R \geq 10 \frac{k}{h} \quad \text{and} \quad \omega_{\varphi^{-1}}(16 \frac{k}{h} \omega_\varphi(\frac{5}{R})) \leq \frac{1}{9}.$$

We proceed by contradiction. Suppose there exists a map $f: G \rightarrow X$ such that $(1/|G|) \sum_{s \in G} \|f(s) - m\| > R \operatorname{Lip}(f)$. We assume without loss of generality that $m = 0$ and $\operatorname{Lip}(f) = 1$. Choose $R_0 > 0$ such that there exists $s_0 \in G$ satisfying

$$|\{x \in G : \|f(s) - f(s_0)\| \leq R_0\}| \geq \frac{3}{4}|G|.$$

We may assume that R_0 attains its infimum under this condition. Since the mean of f is zero, by Lemma A.4, we have

$$R < \frac{1}{|G|} \sum_{s \in G} \|f(s)\| \leq \frac{2}{|G|} \sum_{s \in G} \|f(s) - f(s_0)\| \leq 2R_0 + \frac{k}{h}$$

and hence $R_0 \geq (9/20)R \geq 9k/2h$. Let $\tilde{f}: G \rightarrow X$ be a map given by

$$\tilde{f}(x) = \begin{cases} f(s) & \text{if } \|f(s) - f(s_0)\| \leq R_0 \\ f(s_0) + R_0 \frac{f(s) - f(s_0)}{\|f(s) - f(s_0)\|} & \text{if } \|f(s) - f(s_0)\| > R_0 \end{cases}.$$

It follows that $\tilde{f}$ is a 2-Lipschitz map from G into the closed ball of center $f(s_0)$ and radius R_0 . Let $g: G \rightarrow \ell_1$ be the map given by $g(s) = \varphi((\tilde{f}(s) - f(s_0))/R_0)$. It follows $\operatorname{Lip}(g) \leq \omega_\varphi(2/R_0) \leq \omega_\varphi(5/R)$. Applying lemma A.4 to g , we obtain that

$$\frac{1}{|G|} \sum_{s \in G} \|g(s) - m(g)\| \leq 2\frac{k}{h}\omega_\varphi\left(\frac{5}{R}\right) =: \delta,$$

where $m(g)$ is the mean of g . It follows that for $A = \{s \in G : \|g(s) - m(g)\| \leq 4\delta\}$, we have $|A| \geq (3/4)|G|$. Hence, for $A_1 = A \cap \{s \in G : f(s) = \tilde{f}(s)\}$, we have $|A_1| \geq |G|/2$. Choose $s_1 \in A_1$. Note that for $s \in A_1$, we have $\|g(s) - g(s_1)\| \leq 8\delta$ and hence $\|f(s) - f(s_1)\| \leq \omega_{\varphi^{-1}}(8\delta)R_0 \leq R_0/9$ by the assumption on R . It follows from Lemma A.4 that

$$\frac{1}{|G|} \sum_{x \in G} \|f(s) - f(s_1)\| \leq \frac{1}{9}R_0 + \frac{k}{2h} \leq \frac{2}{9}R_0.$$

Therefore, we get

$$|\{s \in G : \|f(s) - f(s_1)\| \leq \frac{8}{9}R_0\}| \geq \frac{3}{4}|G|,$$

which is in contradiction with the minimality of R_0 . $\square$

REFERENCES

- [Bh] R. Bhatia, *Matrix analysis*. Graduate Texts in Mathematics, **169**. Springer-Verlag, New York, 1997.
- [BL] Y. Benyamini and J. Lindenstrauss, *Geometric nonlinear functional analysis*. Vol. 1. American Mathematical Society Colloquium Publications, **48**. American Mathematical Society, Providence, RI, 2000.
- [Ch] F. Chaatit, *On uniform homeomorphisms of the unit spheres of certain Banach lattices*. Pacific J. Math. **168** (1995), 11–31.

- [Co] A. Connes, *On the cohomology of operator algebras*. J. Funct. Anal. **28** (1978), 248–253.
- [En] P. Enflo, *On a problem of Smirnov*. Ark. Mat. **8** (1969), 107–109.
- [HV] P. de la Harpe and A. Valette, *La propriété (T) de Kazhdan pour les groupes localement compacts*, With an appendix by M. Burger. Astérisque **175** (1989).
- [Jo1] B. E. Johnson, *Cohomology in Banach algebras*. Memoirs of the American Mathematical Society, No. **127**. American Mathematical Society, Providence, R.I., 1972.
- [Jo2] B. E. Johnson, *Approximate diagonals and cohomology of certain annihilator Banach algebras*. Amer. J. Math. **94** (1972), 685–698.
- [Ko] H. Kosaki, *On the continuity of the map $\varphi \rightarrow |\varphi|$ from the predual of a W^* -algebra*, J. Funct. Anal. **59** (1984), 123–131.
- [LLR] N. Linial, E. London, Y. Rabinovich, *The geometry of graphs and some of its algorithmic applications*. Combinatorica **15** (1995), 215–245.
- [Lu] A. Lubotzky, *Discrete groups, expanding graphs and invariant measures*. With an appendix by J. D. Rogawski. Progress in Mathematics, 125. Birkhäuser Verlag, Basel, 1994.
- [Ma] J. Matoušek, *On embedding expanders into ℓ_p spaces*. Israel J. Math. **102** (1997), 189–197.
- [OS] E. Odell and Th. Schlumprecht, *The distortion problem*. Acta Math. **173** (1994), 259–281.
- [Pi1] G. Pisier, *Factorisation de fonctions analytiques à valeurs opérateurs*. C. R. Acad. Sci. Paris Sér. I Math. **307** (1988), 955–960.
- [Pi2] G. Pisier, *On Read’s proof that $B(\ell_1)$ is not amenable*. Seminar Report.
- [PS] R. Powers and E. Størmer, *Free states of the canonical anticommutation relations*, Comm. Math. Phys. **16** (1970) 1–33.
- [Re] C. J. Read, *Relative amenability and the non-amenability of $B(\ell^1)$* . Preprint.
- [Ru] V. Runde, *Lectures on amenability*. Lecture Notes in Mathematics, **1774**. Springer-Verlag, Berlin, 2002.

DEPARTMENT OF MATHEMATICAL SCIENCE, UNIVERSITY OF TOKYO, 153-8914, JAPAN
E-mail address: narutaka@ms.u-tokyo.ac.jp